



Antimicrobial Resistance Profiles of the Two Porcine *Salmonella* Typhimurium Isolates

Kemal METİNER^{1*}, Özden BÜYÜKBABA BORAL², Baran ÇELİK¹, Belgi DİREN SİĞİRCİ¹,
Beren BAŞARAN KAHRAMAN¹, Seyyal AK¹

¹Department of Microbiology, Faculty of Veterinary Medicine, Istanbul University, 34320, Avcılar, Istanbul, Turkey

²Department of Medical Microbiology, Istanbul Faculty of Medicine, Istanbul University, 34390, Çapa, Istanbul, Turkey

***Sorumlu Yazar /
Corresponding Author:**

Kemal METİNER
e-mail: kmetiner@hotmail.com

Geliş Tarihi / Received:
04 November 2015

Kabul Tarihi / Accepted:
27 January 2016

Key Words:
Antimicrobial resistance, ESBL and AmpC
beta-lactamases, pig, *Salmonella*
Typhimurium

Abstract

The aim of the study is to detect the presence of the *Salmonella* species in swine with diarrhea, and to investigate their antimicrobial resistance and extended spectrum beta lactamase (ESBL) and/or AmpC β -lactamase production. For this purpose, stool samples from three commercial pig farms in Istanbul and Tekirdag were collected and processed for *Salmonella* isolation by culture and isolates were identified by biochemical activity tests. *Salmonella* isolates were confirmed by PCR then serotyped. Antimicrobial resistance and ESBL and AmpC production of the isolates were determined according to the Clinical and Laboratory Standards Institute (CLSI) standard. In the study, two hundred and thirty eight stool samples were examined. *Salmonella* spp. were obtained from 2 samples, and the isolation rate was determined as 0.8%. Both of the isolates were defined as *Salmonella enterica* subsp. *enterica* serovar Typhimurium (serotype 1, 4, [5], 12: 1, 2) by serotyping. Both of them were resistant to cefaclor, cloxacillin and lincomycin (100%). Multidrug resistance (resistance ≥ 3 antimicrobials) observed in all isolates. ESBL and AmpC production were not detected in any of the isolates. To our knowledge, this is the first report of the isolation of *S. Typhimurium* in pigs with diarrhea in Turkey. This study also represents the first report of multi-drug resistant *S. Typhimurium* isolates from pig stools in Turkey.

Introduction

Salmonella infections occur in both animals and humans, worldwide. In industrialized countries, non-typhoid *Salmonella* types are an important cause of bacterial gastroenteritis (Van den Brandhof et al., 2003). The most common clinical manifestations of salmonellosis in animals are gastrointestinal disorders; followed by acute septicemia, abortion, arthritis and respiratory disease (Davies, 2004).

Salmonella enterica subsp. *enterica* serovar Typhimurium can cause infections in humans and animals. Most human cases are food borne; however, the bacteria can be transmitted through direct contact and indirectly from the contaminated environment (Rice et al., 2003). *S. Typhimurium* is a common cause of

salmonellosis in many countries (Davies, 2004; Mmolawa et al., 2002) and it was reported that the predominant serotype in Europe (van Pelt et al., 2003). In addition, it was indicated that *S. Typhimurium* strains were the most important serotypes that frequently causes food-related and widespread infections in Turkey (Erdem et al., 2005).

The infected pigs with *Salmonella* types, shed the bacteria through faeces that was caused by contamination of slaughterhouses and might be created a potential health risk for humans (Swanenburg et al., 2001).

It was a known fact that, infections caused by antimicrobial resistant bacteria in humans and animals has been increasing lately (Arias and Murray, 2015;

Bagcigil et al., 2007; Yilmaz et al., 2015). It was suggested that the increase usage of cephalosporins in human and veterinary medicine might be the main cause of extended-spectrum β -lactamase (ESBL) and AmpC β -lactamase producing bacteria colonization. β -lactamase enzymes are mainly plays an important role in the Enterobacteriaceae, and β -lactamase resistance could be transferred among other bacteria via plasmids (Li et al., 2007).

Presence of multi-drug resistance (resistance to ≥ 3 antimicrobials) were created a potential risk for the treatment of the infections and restrain the drug choice (Van Boxtael et al., 2012). The drug resistance rate, varies between different serotypes, some *S. Typhimurium* strains have been reported resistant to 10 or more antimicrobial agents (Poppe et al., 2002).

The aim of the study is to detect the presence of the *Salmonella* species in swine, and to investigate their antimicrobial resistance and ESBL and/or AmpC β -lactamase production.

Materials and Methods

Stool samples were taken from 238 pigs with diarrhea from three different farms in Çorlu (Tekirdağ), Ayazma and Arnavutköy (Istanbul). The stool samples were collected into sterile sample tubes and transported to the laboratory in the cold chain.

The culture was conducted to the standard of *Salmonella* isolation from stool samples. Biochemical tests were carried out by Api 20 E (Bio Merieux, France) (Ertaş et al., 2014; Holt et al., 1994; Türkyılmaz et al., 2007). Confirmation of the genus *Salmonella* was performed by genus-specific PCR according to Rahn et al. (1992), and Shanmugasamy et al. (2011). *S. enteritidis* strain (KUEN 349) was used as the positive control. The serotyping of the isolates was carried out on the Veterinary Control and Research Institute, *Salmonella* Reference Laboratory, Etlik, Ankara.

Antimicrobial resistance of the isolates was determined by the Kirby-Bauer disc diffusion method according to the Clinical Laboratory Standards Institute (CLSI) manual (2013). The tested antibiotics are; ampicillin (AMP 10 μ g), amoxicillin (AML 10 μ g), amoxicillin/clavulonic acid (AMC 10 μ g), cefaclor (CEC 30 μ g), cephalexin (CL 30 μ g), cefoperazone (CFP 75 μ g), ceftiofur (CFT 30 μ g), chloramphenicol (C 30 μ g), ciprofloxacin (CIP 5 μ g), cloxacillin (OB 6 μ g), enrofloxacin (ENR 5 μ g), gentamycin (GN 5 μ g), kanamycin (K 30 μ g), lincomycin (MY 2 μ g), nalidixic acid

(NA 30 μ g), nitrofurantoin (F 300 μ g), streptomycin (S 10 μ g), sulbactam/ampicillim (SAM 20 μ g) and sulfamethoxazole/trimethoprim (SXT 25 μ g).

ESBL screening and confirmatory tests of the isolates were evaluated according to the guidelines of the CLSI by Kirby-Bauer disc diffusion tests. The susceptibility was tested against cefotaxime (30 μ g), ceftriaxone (30 μ g), ceftazidim (30 μ g), cefpodoxim (10 μ g), aztreonam (30 μ g) (HiMedia) in screening tests. The diameter of the zone of inhibition was measured and interpreted according to ESBL screening test limits in CLSI (2013). The production of phenotypic AmpC β -lactamase was determined by resistance to ceftiofur (30 μ g) (≤ 14 mm) and susceptibility to cefepime (30 μ g) (≥ 18 mm) (CLSI, 2013; EFSA, 2011; Giske et al., 2012).

Results

Salmonella spp. isolated in 2 of the 238 samples, the isolation rate was determined as 0.8%. Both of the isolates were obtained from the samples collected to the same source. Additionally, isolates were verified as *Salmonella* by PCR (Figure 1) and *Salmonella enterica* subsp. *enterica* serovar Typhimurium (serotype 1, 4, [5], 12: I: 1, 2) by serotyping.

Multiple antibiotic resistance determined in the both of the isolates. Antimicrobial resistance patterns of the isolates were shown in the Table 1. The productions of phenotypic ESBL and AmpC β -lactamase have not been detected in any of the isolates. The results of the screening tests were shown in Table 2.




Figure 1. Agarose gel electrophoresis image of PCR products.

Lane 1; 100 bp molecular weight marker, lane 2; positive sample 1, lane 3; positive sample 2, lane 4; positive control, lane 5; negative control.

Table 1. Antimicrobial resistance patterns of the isolates.

Resistance Pattern			
Isolate 1	CEC	OB	MY
Isolate 2	CEC	OB	MY

CEC: Cefaclor, OB: Cloxacillin, MY: Lincomycin

Table 2. Phenotypic ESBL and AmpC β -lactamase assessments of the isolates.

ESBL Screening Test					Result	AmpC Definition Test			Result
Inhibition Zone Diameter (mm)				Inhibition Zone Diameter (mm)					
CTX	CTR	CAZ	CPD	CTX		CX	CAZ		
Isolate 1	26	24	24	23	-	26	22	22	-
Isolate 2	28	29	24	22	-	26	21	21	-

CTX: Cefotaxime, CTR: Ceftriaxone, CAZ: Ceftazidime, CPD: Cefpodoxime, CX: Cefoxitin, -: Negative

Discussion

Salmonella infections may observe in all of the domestic animals and humans. The young ones, pregnant and lactating animals are the most susceptible proportion of the populations. Enteric disease is the most common clinical manifestation, but a wide range of clinical signs, which include acute septicaemia, abortion, arthritis and respiratory disease, may be seen. (Baggesen et al., 2000; Boyen et al., 2008).

There were a limited number of studies carried out on bacterial diseases in pigs in Turkey (Bağcıgil et al., 2013; Metiner et al., 2007; Metiner et al., 2013). Likewise, during the literature search at the planning stage of the current study, has not been encountered any sources for the investigation of Salmonella shedding in pigs raised in Turkey. However, there have been some studies on colonization of *Salmonella* spp. in cattle, sheep and poultry (Aydın et al., 2001; Genç, 2002; Genç and Otlı, 2005).

In the world wide, *Salmonella* spp. isolation rate from swine were reported between 4.3% and 27.5% (Bolton et al., 2013; Li et al., 2013; Vieira-Pinto et al., 2006; White et al., 2004). In the current study, isolation rate was much lower than the rates of the previous studies. It was thought that the low isolation rate of *Salmonella* types could be based on limited number of swine population in Turkey.

The multidrug-resistance of the *Salmonella* types reduces the therapeutic options in cases of invasive infections in both human and animals. Rajic et al. (2004), reported that was observed of resistance for

chloramphenicol (4.7%), ampicillin (7.8%), kanamycin (11.8%), sulphamethoxazole (21.1%), streptomycin (25.5%) and tetracycline (38.8%). Wright et al. (2005), was reported that multi-drug resistant *S. Typhimurium* was isolated from an epidemic. Sisak et al. (2006), indicated that the multi-drug resistance in pig isolates, and the isolates were resistant to amoxicillin/clavulonic acid, sulphamethoxazole/trimethoprim, nalidixic acid and enrofloxacin. De Jong et al. (2014), reported that the rate of multi-drug resistance in pig as 39.7%. In the current study, two *Salmonella* isolates had multi-drug resistance with cefaclor, cloxacillin and lincomycin.

It was emphasized that the widespread use of the antibiotics in animals may lead to the increase of the colonization of resistant bacteria such as ESBL and AmpC β -lactamase producers and the resistant strains may spread to the society (Davies, 2004). It was reported that ESBL and AmpC β -lactamases can be transmitted between the bacteria with their transferable genetic elements and clinical over use of the β -lactam antibiotics may cause to a significant public health risk (Frye and Fedorka-Cray, 2007; Li et al., 2007). In a study that was carried out in the USA, indicated that the ceftiofur resistance in *Salmonella* strains were 9.8% in cats, 19.2% in horses and 20.8% in dogs (Frye and Fedorka-Cray, 2007). De Jong et al. (2014) reported that two of 128 *S. Typhimurium* isolates were producing ESBL while two of them were producing AmpC β -lactamase. In the current study, any of the isolates were produced ESBL and AmpC β -lactamase.

In conclusion, in the current study, isolation rate was much lower than the rates of the previous studies. It

was thought that the reason for this result, there were a limited swine population in Turkey, compared to other countries. On the other hand, both of the isolates (100%) had multi-drug resistance to the antimicrobials. When the transmission risk rather low, the results were suggested that if the infections caused by this species happened, it would be difficult to treat.

To our knowledge, this is the first report of the isolation of *S. Typhimurium* in pigs with diarrhea in Turkey. This study also represents the first report of multi-drug resistant *S. Typhimurium* isolates from pigs' stools in Turkey.

REFERENCES

- Arias, C.A., Murray, B.E., 2015. A new antibiotic and the evolution of resistance. *New England Journal of Medicine* 372, 1168-1170.
- Aydın, F., Umur, Ç., Gökçe, G., Genç, O., Güler, M.A., 2001. The isolation and identification of bacteria and parasites from diarrhoeic calves in Kars District. *Journal of the Faculty of Veterinary Medicine, Kafkas University* 7, 7-14.
- Bağcigil, A.F., İkiz, S., Metiner, K., Özgür, N.Y., Ak, S., Ilgaz, A.A., 2013. Isolation of *Streptococcus* species from the tonsils of slaughtered pigs. *Turkish Journal of Veterinary and Animal Sciences* 37, 94-96.
- Bagcigil, F.A., Moodley, A., Baptiste, K.E., Jensen, V.F., Guardabassi, L., 2007. Occurrence, species distribution, antimicrobial resistance and clonality of methicillin- and erythromycin-resistant staphylococci in the nasal cavity of domestic animals. *Veterinary Microbiology* 121, 307-315.
- Baggesen, D.L., Sandvang, D., Aarestrup, F.M., 2000. Characterization of *Salmonella enterica* *Salmonella enterica* serovar Typhimurium DT104 isolated from Denmark and comparison with isolates from Europe and the United States. *Journal of Clinical Microbiology* 38, 1581-1586.
- Bolton, D.J., Ivory, C., McDowell, D., 2013. A study of *Salmonella* in pigs from birth to carcass: serotypes, genotypes, antibiotic resistance and virulence profiles. *International Journal of Food Microbiology* 160, 298-303.
- Boyen, F., Haesebrouck, F., Maes, D., van Immerseel, F., Ducatelle, R., Pasmans, F., 2008. Non-typhoidal *Salmonella* infections in pigs: A closer look at epidemiology, pathogenesis and control. *Veterinary Microbiology* 130, 1-19.
- CLSI (Clinical and Laboratory Standards Institute), 2013. M100-S23-performance standards for antimicrobial susceptibility testing; twenty-third informational supplement. CLSI, Wayne, Pennsylvania.
- Davies, R., 2004. *Salmonellosis*. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. OIE, World Organization for Animal Health, 1018-1032.
- de Jong, A., Smet, A., Ludwig, C., Stephan, B., De Graef, E., Vanrobaeys, M., Haesebrouck, F., 2014. Antimicrobial susceptibility of *Salmonella* isolates from healthy pigs and chickens (2008–2011). *Veterinary Microbiology* 171, 298-306.
- EFSA (European Food Safety Authority), 2011. Scientific opinion on the public health risks of bacterial strains producing extended-spectrum β -lactamases and/or ampc β -lactamases in food and food-producing animals. EFSA Panel on Biological Hazards (BIOHAZ). *EFSA Journal* 9, 2322, 1-95.
- Erdem, B., Ercis, S., Hascelik, G., Gur, D., Gedikoglu, S., Aysev, A.D., Sumerkan, B., Tatman-Otkun, M., Tuncer, I., 2005. Antimicrobial resistance patterns and serotype distribution among *Salmonella enterica* strains in Turkey, 2000-2002. *European Journal of Clinical Microbiology and Infectious Diseases* 24, 220-225.
- Ertaş, N., Al, S., Karadal, F., Gönülalan, Z., 2014. Kayseri ilinde Satışa Sunulan Manda Yoğurtlarının Mikrobiyolojik Kalitesi. *İstanbul Üniversitesi Veteriner Fakültesi Dergisi* 40, 83-89.
- Frye, J.G., Fedorka-Cray, P.J., 2007. Prevalence, distribution and characterisation of ceftiofur resistance in *Salmonella enterica* isolated from animals in the USA from 1999 to 2003. *International Journal of Antimicrobial Agents* 30, 134-142.
- Genç, O., 2002. The isolation, identification and serotyping of *Salmonella* isolated from domestic animals in Kars district. *Journal of the Faculty of Veterinary Medicine, Kafkas University* 8, 23-30.
- Genç, O., Otlı, S., 2005. *Salmonella* isolations from different avian species in Kars district of Turkey. *The Journal of the Faculty of Veterinary Medicine, Kafkas University* 11, 25-27.
- Giske, C.G., Martinez-Martinez, L., Cantón, R., Stefani, S., Skov, R., Glupczynski, Y., Poland, M.G., 2012. EUCAST Guidelines for Detection of Resistance Mechanisms and Specific Resistances of Clinical and/or Epidemiological Importance. *European Committee on Antimicrobial Susceptibility Testing*, 1-47.
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T., Williams, S.T., 1994. *Bergey's Manual of Determinative Bacteriology*, Ninth Edition. Lippincott Williams and Wilkins, USA.
- Li, R., Lai, J., Wang, Y., Liu, S., Li, Y., Liu, K., Shen, J., Wu, C., 2013. Prevalence and characterization of *Salmonella* species isolated from pigs, ducks and chickens in Sichuan Province, China. *International Journal of Food Microbiology* 163, 14-18.
- Li, X.Z., Mehrotra, M., Ghimire, S., Adewoye, L., 2007. β -Lactam resistance and β -lactamases in bacteria of animal origin. *Veterinary Microbiology* 121, 197-214.
- Metiner, K., Ak, S., 2007. Presence and seroprevalence of *Actinobacillus pleuropneumoniae* in pigs in Turkey. *Acta Veterinaria Brno* 76, 237-244.

- Metiner, K., Anđ Küçüker, M., Büyükbaba Boral, Ö., Anđ, Ö., 2013.** First isolation of *Enterococcus* strains in pig faeces in Turkey and determination of antibiotic susceptibilities. *Acta Veterinaria Brno* 82, 231-235.
- Mmolawa, P.T., Willmore, R., Thomas, C.J., Heuzenroeder, M.W., 2002.** Temperate phages in *Salmonella enterica* serovar Typhimurium: Implications for epidemiology. *International Journal of Medical Microbiology* 291, 633-644.
- Poppe, C., Ziebell, K., Martin, L., Allen, K., 2004.** Diversity in Antimicrobial Resistance and Other Characteristics Among *Salmonella* Typhimurium DT104 Isolates. *Microbial Drug Resistance* 8, 107-122.
- Rahn, K., De Grandis, S.A., Clarke, R.C., McEwen, S.A., Galán, J.E., Ginocchio, C., Curtiss III, R., Gyles, C.L., 1992.** Amplification of an *invA* gene sequence of *Salmonella* Typhimurium by polymerase chain reaction as a specific method of detection of *Salmonella*. *Molecular and Cellular Probes* 6, 271-279.
- Rajić, A., McFall, M.E., Deckert, A.E., Reid-Smith, R., Manninen, K., Poppe, C., Dewey, C.E., McEwen, S.A., 2004.** Antimicrobial resistance of *Salmonella* isolated from finishing swine and the environment of 60 Alberta swine farms. *Veterinary Microbiology* 104, 189-196.
- Rice, D.H., Hancock, D.D., Roozen, P.M., Szymanski, M.H., Scheenstra, B.C., Cady, K.M., Besser, T.E., Chudek, P.A., 2003.** Household Contamination with *Salmonella enterica*. *Emerging Infectious Diseases* 9, 120-122.
- Shanmugasamy, M., Velayutham, T., Rajeswar, J., 2011.** *InvA* gene specific PCR for detection of *Salmonella* from broilers. *Veterinary World* 4, 562-564.
- Sisak, F., Havlickova, H., Hradecka, H., Rychlik, I., Kolackova, I., Karpiskova, R., 2006.** Antibiotic resistance of *Salmonella* spp. isolates from pigs in the Czech Republic. *Veterinari Medicina* 51, 303-310.
- Swanenburg, M., Berends, B.R., Urlings, H.A.P., Snijders, J.M.A., van Knapen, F., 2001.** Epidemiological investigations into the sources of *Salmonella* contamination of pork. *Berliner und Münchener Tierärztliche Wochenschrift* 114, 356-359.
- Türkyılmaz, S., Savaşan, S., Kirkan, Ş., Kaya, O., 2007.** Detection of *Salmonella* Enteritidis infections in poultry by using bacteriological and serological methods. *Journal of the Faculty of Veterinary Medicine Istanbul University* 33, 23-33.
- Van Boxtael, S., Dierick, K., Van Huffel, X., Uyttendaele, M., Berkvens, D., Herman, L., Bertrand, S., Wildemaue, C., Catry, B., Butaye, P., Imberechts, H., 2012.** Comparison of antimicrobial resistance patterns and phage types of *Salmonella* Typhimurium isolated from pigs, pork and humans in Belgium between 2001 and 2006. *Food Research International* 45, 913-918.
- Van den Brandhof, W.E., van Pelt, W., Wagenaar, J.A., van Duynhoven, Y.T.H.P., 2003.** Study on the sources of human campylobacteriosis and salmonellosis in the Netherlands. *Infectieziekten Bulletin* 14, 132-135.
- Van Pelt, W., de Wit, M.A.S., Wannet, W.J.B., Ligtoet, E.J.J., Widdowson, M.A., van Duynhoven, Y.T.H.P., 2003.** Laboratory surveillance of bacterial gastroenteric pathogens in The Netherlands, 1991-2001. *Epidemiology and Infection* 130, 431-441.
- Vieira-Pinto, M., Tenreiro, R., Martins, C., 2006.** Unveiling contamination sources and dissemination routes of *Salmonella* sp. in pigs at a Portuguese slaughterhouse through macrorestriction profiling by pulsed-field gel electrophoresis. *International Journal of Food Microbiology* 110, 77-84.
- White, D.G., Zhao, S., Singh, R., McDermoot, P.F., 2004.** Antimicrobial Resistance Among Gram-Negative Foodborne Bacterial Pathogens Associated with Foods of Animal Origin. *Foodborne Pathogens and Disease* 1, 137-152.
- Wright, J.G., Tengelsen, L.A., Smith, K.E., Bender, J.B., Frank, R.K., Grendon, J.H., Rice, D.H., Thiessen, A.M.B., Gilbertson, C.J., Sivapalasingam, S., Barret, T.J., Besser, T.E., Hancock, D.D., Angulo, F.J., 2005.** Multidrug-resistant *Salmonella* Typhimurium in Four Animal Facilities. *Emerging Infectious Diseases* 11, 1235-1241.
- Yılmaz, M.A., Aslantaş, Ö., Özer, B., Yılmaz, E.Ş., 2015.** Nasal Carriage of Methicillin-Resistant Coagulase Negative Staphylococci (MR-CoNS) Among Veterinarians and Veterinary Students. *Journal of the Faculty of Veterinary Medicine Istanbul University* 41, 69-78.