

# Influence of biologically active and phytogetic feed supplements on the productiveness in growing and fattening pigs

\*Gergana Yordanova<sup>1</sup>, Radka Nedeva<sup>1</sup>

<sup>1</sup>Agricultural Institute – Shumen

\*E-mail: gergana22241@abv.bg

**Citation:** Yordanova, G. & Nedeva, R. (2020). Influence of biologically active and phytogetic feed supplements on the productiveness in growing and fattening pigs. *Zhivotnovadni Nauki*, 57(5), 41-45 (Bg).

## Abstract

Phytogetic feed supplements are plant products used in animal nutrition in order to improve productivity. This class of feed has recently gained interest, especially since the EU ban on antibiotic feed supplements in 2006. Phytogetics is a fairly new class of feed supplements, with limited research on their effects and application, as well as studies on their botanical origin, processing and composition. The literature analysis indicated that herbal and plant extracts are one of the alternatives for growth promoters as their dosage, type and combinations of application will continue to be a subject of study for nutritional science.

This article considers researches on the influence of herbs and herbal extracts, activated charcoal and *Spirulina plantesis* in combined feeds, the productiveness and health of growing and fattening pigs.

**Key words:** nutrition, biologically active supplements, pigs, health, productiveness

The developmental directions of pig nutrition science in recent years are related to the use of biologically active and phytogetic feed supplements, which provide improvement of the microflora of digestion and absorption of nutrients and maintaining the health status of animals.

Phytogetic feed supplements are plant products used in animal nutrition in order to improve productivity. This class of feed has recently gained interest, especially since the EU ban on antibiotic feed supplements in 2006. Phytogetics is a fairly new class of feed supplements, which do not include organic acids or probiotics that have been widely studied in animal nutrition. Studies about phytogetic supplements, their effects and application are limited, as well as studies on their botanical origin, processing and composition (Windisch et al., 2007).

Studies by a number of authors (Namkung et al., 2004, Demir et al., 2005, Jamroz et al., 2006, Nofrarias et al., 2006, Cho et al., 2006, Stoni et al., 2006, Oetting et al., 2006) have been aimed at establishing morphological changes in the gastrointestinal tissues in pigs and birds due to the use of phytogetic feed supplements. Studies show both an increase and no change or decrease in the length of the intestinal villi in pigs and birds. The authors substantiated the claims by supporting the hypothesis that these supplements may stabilize nutrition.

In their review, Windisch et al. (2007) described possible antioxidant, antimicrobial and growth effects, as well as the effects in animals from the use of phytogetic supplements associated with improved consumption of compound feeds. Some researchers have shown the poten-

tial of phytogenic supplements to reduce intestinal pathogenic pressure, in terms of antimicrobial activity. According to some studies, they have the ability to modulate microbial colonies, fermentation and nutrient digestibility. Some isolated observations appear to support improvement in digestive enzyme activity and absorption capacity through the use of phytogenic compounds.

The ban on antibiotic application (2006) as growth stimulants has led to the search for alternative biological supplements that ensure the preservation and health of animals, as well as their productivity (Ivanova et al., 2010; Ivanova-Peneva et al., 2006; Borovan, 2004; Huang et al., 2011). Plant extracts have been used in animal nutrition especially because of their antimicrobial (Namkung et al., 2004; Costa et al., 2007), anti-inflammatory and antioxidant (Liu et al., 2008) and antiparasitic properties (Magi et al., 2006). It has been established that some natural feed products can have a prophylactic effect against a number of intestinal diseases due to their immunological and pharmacological action (Laine et al., 2008).

The effect of adding herbs in pig farming, especially in young pigs, has mainly signified, on the one hand, the improvement in health and on the other hand the improvement of some productive indicators – growth, consumption and utilization of feed.

In the studies of Ivanova (2012) the combination of the *Organum vulgare* and *Potentilla Erecta Raus* herbs did not have an effect on feed consumption, average daily gain and feed utilization in grower pigs. However, the addition of this combination of herbs, before and after weaning has a positive influence on post-weaning diarrhea prevention.

The addition of herbs to the water or feed of suckling pigs improved their growth and health state (Ivanova-Peneva et al., 2006).

The use of *Organum vulgare* and *Potentilla Erecta Raus* in combined feeds for suckling pigs has been studied by Ivanova-Peneva and others (2010). The authors established an improvement in growth (by 11%,  $p \leq 0.01$ ) for the suckling period (1–34 days) and live weight at weaning

(7.500 kg vs. 6.883 kg and 6.944 kg) due to the added herbs. The conclusion of the authors was that the herbs *Organum vulgare* and *Potentilla Erecta Raus* have a therapeutic effect and can be successfully used in the rearing of suckling pigs for prophylactic and therapeutic purposes.

The effect of the addition of a mixture of plant extracts, including *Rehmannia glutingsa*, *Angelica gigas*, *Discorea japonica*, *Glucurrrhiza uralensis*, *Glycyrrhiza uralensis*, *Schisan-dra chinensis* and *Ligusticum jeholense*, on the productivity of some blood indicators was studied by Lee Paik (2007). Gain and feed intake were significantly higher in the groups receiving herbal mixture, and no difference in dosages was found. The addition of the herbal mixture decreased the content of *E. coli*, while those of *Lactobacilli* were increased.

The addition of an herbal mixture (2.04 and 3.0%) to the rations of fattened pigs in studies done by Paschma et al. (2010) reduced ammonium emissions over three test seasons (summer, autumn and winter).

The inclusion of Chinese herbs (1.0% and 0.5%) in studies done by Chen et al. (2011) increased live weight by 9.89% and 4.42% and reduced feed consumption by 7.07% and 4.49%. The authors also reported a positive effect of the added Chinese herbs on liver and kidney functions.

A number of authors (Castillo et al., 2006; Alakomi et al., 2006; Santoro et al., 2007; Moon et al., 2006) found that the beneficial effect of herbs and plant extracts on the gastrointestinal microflora was mainly related to the inhibitory effect on various intestinal parasites and bacteria.

Kanev et al. (2002) studied the effect of Ropadiar (essential oils of *Organum vulgare*, thyme and sage) on the growth of suckling pigs. The authors found that the use of Ropadiar at a dose of 0.05% in the formula of suckling pigs from 7 to 35 days of age provided higher feed, nutrients and energy consumption and higher average daily gain.

In her thesis, Zapryanova-Boeva (2011) tested the biologically active plant supplement Vemo-Herb (extracts of Chicory, Smoke tree and tansy)

on the productive traits of growing and fattening pigs, sows and their offspring. The author found that the supplement improved growth rate by 29% compared to the animals that hadn't been fed with growth stimulant and up to 18.8% compared to groups with nutritional antibiotic supplementation. The addition of VemoHerb increased the gain of fattened pigs by 9 to 18%. In sows, the addition of herbal extracts had increased feed consumption, live weight of piglets at birth and their gain during the suckling period.

The addition of conventional foods with *Spirulina platensis* seems to be a promising trend due to the nutritional value of this type of algae, which is known to be a rich source of protein, fatty acids, fiber, essential vitamins and minerals. It also has functional properties due to the presence of antioxidants, omega 3, antiviral and anti-cancer components. When *Spirulina platensis* is added to processed cheese, the content of protein, ash, fiber, selenium, zinc, iron, magnesium and potassium is increased in order to increase the nutritional and health value of the final product. There has been an increase in the activity of antioxidants, improvement of organoleptic, physical and rheological properties, which has led to the conclusion that the supplement has excellent nutritional, functional and health properties that can be used as a food product (Tohamy and al., 2019).

Natural plant charcoal is a feed additive, which we tested in weaned pigs. The supplement is obtained from specially selected French oak by carbonization. The product has a very good absorption capacity and captures enterotoxins produced by harmful bacteria and mycotoxins from feed, as well as some pathogenic bacteria such as clostridia, salmonella and *E. coli*.

Activated charcoal is a fine black powder that is obtained from the decomposition of various organic materials, which is then exposed to oxidizing gases at high temperatures to activate and increase the surface area (Clegg and Hope, 1999).

Struhsaker et al. (1997) stated that activated charcoal adsorbs a wide range of compounds. Adsorption therapy with activated charcoal as an indigestible carrier has been one of the im-

portant methods to prevent the ingestion of toxic substances or harmful substances formed in the gastrointestinal tract (McLennan and Amos, 1989; McKenzie, 1991; Jindal et al., 1994).

Chu et al. (2013) found that the inclusion of bamboo charcoal in the rations of fattened pigs increased productivity and feeding efficiency by reducing gaseous emissions and harmful microflora in faeces. The authors concluded that the charcoal used has been shown to reduce stress as reported by lower cortisol levels and increased serum Ig G concentrations.

Some researchers have studied charcoal and activated charcoal as a feed supplement. Charcoal affected the growth and carcass characteristics of pigs for fattening (Hwang, 1995) and was used as a feed additive for the production of high quality meat (Kim, 1990). Activated charcoal also affected meat quality and pork storage (Lee et al., 2011).

Activated charcoal supplementation in rations of fattening pigs has not been tested locally. In our research with grower pigs (Nedeva and Yordanova, 2013) we established that the addition of plant charcoal (3 kg/t feed) during a 14, 21, 35 and 49-day period after weaning increased growth intensity with 17.65–24.81% and decreased feed consumption and nutrition with 5.47–18.17%. In the conditions of our experiment the added plant supplement has successfully decreased the number of digestive disorders.

The literature analysis indicated that herbal and plant extracts are one of the alternatives for growth promoters as their dosage, type and combinations of application will continue to be a subject of study for nutritional science.

Biodiversity in nature, including microorganisms, terrestrial plants, seaweed and marine organisms, offers a valuable source of new bioactive substances. In reference to this we are monitoring the progress of development in immune mechanisms, through which the dynamic interaction of the intestinal microbiota and its host usually favours the homeostatic, symbiotic relationship and the way macroalgal bioactive substances are fed in both maternal and piglet rations can be used to support this symbiotic relationship (Sweeney et. al., 2016).

In summary, the reasonable use of antibiotics and the continuous development of alternatives are needed for ensuring the long-term sustainable development of livestock breeding.

At the same time, we need to strengthen surveillance and law enforcement to control antibiotic resistance and food chain residues within established safe levels. In addition, we need to improve the management of animal nutrition and production hygiene. Exploring alternatives to antibiotics will be a long process. In addition to research and development of new effective and safe alternatives, we need to step up research into the effects of the combined use of antibiotics and their alternatives, aimed at maintaining a healthy agricultural economy and preserving powerful antibiotics for effective human therapy.

## References

- Alakomi, H. L., Paananen, A., Suihko, M. L., Helander, I. M., & Saarela, M.** (2006). Weakening effect of cell permeabilizers on gram-negative bacteria causing biodeterioration. *Applied and environmental microbiology*, 72(7), 4695-4703.
- Borovan, L.** (2004). Plant alkaloids enhance performance of animals and improve the utilizability of amino acids. *Krmivarstvi*, 6, 36-37.
- Castillo, M., Martín-Orúe, S. M., Roca, M., Manzanilla, E. G., Badiola, I., Perez, J. F., & Gasa, J.** (2006). The response of gastrointestinal microbiota to avilamycin, butyrate, and plant extracts in early-weaned pigs. *Journal of Animal Science*, 84(10), 2725-2734.
- Chen, Z., Guo, S., Kong, D., & Wei, G.** (2011). Effects of Chinese herb additives on growth performance and blood biochemical index in fattening pigs. *Chinese Journal of Veterinary Science*, 31(10), 1523-1525.
- Cho, J. H., Chen, Y. J., Min, B. J., Kim, H. J., Kwon, O. S., Shon, K. S., Kim, I. H., Kim, S., & Asamer, A.** (2006). Effects of essential oils supplementation on growth performance, IgG concentration and fecal noxious gas concentration of weaned pigs. *Asian-Australasian Journal of Animal Sciences*, 19(1), 80-85.
- Chu, G. M., Kim, J. H., Kim, H. Y., Ha, J. H., Jung, M. S., Song, Y., Cho, J., Lee, S., Ibrahim, R., Lee, S. S., & Song, Y. M.** (2013). Effects of bamboo charcoal on the growth performance, blood characteristics and noxious gas emission in fattening pigs. *Journal of Applied Animal Research*, 41(1), 48-55.
- Clegg, T., & Hope, K.** (1999). The first line of response for people who self-poison: exploring the options for gut decontamination. *Journal of advanced nursing*, 30(6), 1360-1367.
- Costa, L. B., Tse, M. L. P., & Miyada, V. S.** (2007). Herbal extracts as alternatives to antimicrobial growth promoters for weanling pigs. *Revista Brasileira de Zootecnia*, 36(3), 589-595.
- Demir, E., Sarica, S., Ozcan, M. A., & Suicmez, M.** (2005). The use of natural feed additives as alternatives to an antibiotic growth promoter in broiler diets. *Archiv fur Geflugelkunde*, 69(3), 110-116.
- Huang, R. H., Qiu, X. S., Shi, F. X., Hughes, C. L., Lu, Z. F., & Zhu, W. Y.** (2011). Effects of dietary allicin on health and growth performance of weanling piglets and reduction in attractiveness of faeces to flies. *Animal: an international journal of animal bioscience*, 5(2), 304-311.
- Hwang, M. J.** (1995). Effects of activated carbon on the growth rate, feed efficiency, and carcass characteristics in pigs. *Ph.D. Thesis*, Seoul, Korea: KonKuk University.
- Ivanova-Peneva, S.** (2012). Application of herbs for the prevention and treatment of after weaning diarrhoea, stimulation of growth and improving welfare of growing pigs. *Bulgarian Journal of animal husbandry*, 6, 34-42; (bg)
- Ivanova-Peneva, S., Gineva, E., & Nedeva, R.** (2010). Effect of *Origanum vulgare* and *Potentilla Erecta* Raus on the productiveness and health condition of suckling piglets. *Zhivotnov'dni Nauki*, 47(4), 32-41.
- Ivanova-Peneva, S. G., Nedeva, R. D., & Kirov, M. S.** (2006). Preliminary studies on the effect of herbs on the growth and health of suckling piglets. *Proceeding of Joint Organic congress "Organic Farming and European Rural Development"*, Odense, Denmark, 558-559.
- Jamroz, D., Wartecki, T., Houszka, M., & Kamel, C.** (2006). Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *Journal of Animal Physiology and Animal Nutrition*, 90(5-6), 255-268.
- Jindal, N., Mahipal, S. K., & Mahajan, N. K.** (1994). Toxicity of aflatoxin B1 in broiler chicks and its reduction by activated charcoal. *Research in Veterinary Science*, 56(1), 37-40.
- Kanev, D., Petrov, P., & Nedeva, R.** (2002). Effect of Ropadiar used in suckler pig mixtures. *Journal of Mountain Agriculture on the Balkans*, том V, 3, 226-234. (bg)
- Kim, D. H.** (1990). A study of utilizability as feed additives for ground charcoal made of condensed sawdust on the broiler production. *Graduate School of KonKuk University*. [in Korean].
- Laine, T. M., Lyytikäinen, T., Yliaho, M., & Anttila, M.** (2008). Risk factors for post-weaning diarrhoea

on piglet producing farms in Finland. *Acta Veterinaria Scandinavica*, 50(1), 21.

**Lee, W. S., & Paik, I. K.** (2007). Effects of Herb Mix® supplementation on the performance of weanling pigs. *Journal of Animal Science and Technology*, 49(3), 321-328.

**Lee, J. J., Park, S. H., Jung, D. S., Choi, Y. I., & Choi, J. S.** (2011). Meat quality and storage characteristics of finishing pigs by feeding stevia and charcoal. *Korean Journal of Food Science of Animal Resources*, 31(2), 296-303.

**Liu, G. M., Wei, Y., Wang, Z. S., Wu, D., Zhou, A. G., & Liu, G. L.** (2008). Effects of herbal extract supplementation on growth performance and insulin-like growth factor (IGF)-I system in finishing pigs. *Journal of Animal and Feed Sciences*, 17(4), 538-547.

**Mägi, E., Järvis, T., & Miller, I.** (2006). Effects of different plant products against pig mange mites. *Acta Veterinaria Brno*, 75(2), 283-287.

**McKenzie, R. A.** (1991). Bentonite as therapy for Lantana camara poisoning of cattle. *Australian Veterinary Journal*, 68(4), 146-148.

**McLennan, M. W., & Amos, M. L.** (1989). Treatment of lantana poisoning in cattle. *Australian veterinary journal*, 66(3), 93-94.

**Moon, T., Wilkinson, J. M., & Cavanagh, H. M.** (2006). Antiparasitic activity of two Lavandula essential oils against Giardia duodenalis, Trichomonas vaginalis and Hexamita inflata. *Parasitology research*, 99(6), 722-728.

**Namkung, H., Li J. Gong, M., Yu, H., Cottrill, M., & De Lange, C. F. M.** (2004). Impact of feeding blends of organic acids and herbal extracts on growth performance, gut microbiota and digestive function in newly weaned pigs. *Canadian journal of animal science*, 84(4), 697-704.

**Nedeva, R., & Yordanova, G.** (2013). Effect of the addition of carbovet on the productivity and health in growing pigs. *Bulgarian Journal of Animal Husbandry*, 2, 14-18 (bg)

**Nofrarias, M., Manzanilla, E. G., Pujols, J., Gibert, X., Majo, N., Segalés, J., & Gasa, J.** (2006). Effects of spray-dried porcine plasma and plant extracts on intesti-

nal morphology and on leukocyte cell subsets of weaned pigs. *Journal of Animal Science*, 84(10), 2735-2742.

**Oetting, L. L., Utiyama, C. E., Giani, P. A., Ruiz, U. D. S., & Miyada, V. S.** (2006). Effects of herbal extracts and antimicrobials on apparent digestibility, performance, organs morphometry and intestinal histology of weanling pigs. *Revista Brasileira de Zootecnia-Brazilian Journal of Animal Science*, 35(4), 1389-1397.

**Paschma, J., Kaczor, A., & Paraponiak, P.** (2010). Evaluation of the possibility of improving the breeding environment of pigs receiving dietary herb mixture. *Roczniki Naukowe Zootechniki*, 37(2), 179-186.

**Santoro, G. F., das Graças Cardoso, M., Guimarães, L. G. L., Salgado, A. P. S., Menna-Barreto, R. F., & Soares, M. J.** (2007). Effect of oregano (Origanum vulgare L.) and thyme (Thymus vulgaris L.) essential oils on Trypanosoma cruzi (Protozoa: Kinetoplastida) growth and ultrastructure. *Parasitology research*, 100(4), 783-790.

**Stoni, A., Zitterl-Egelseer, K., Kroismayr, A., Wetscherek, W., & Windisch, W.** (2006, March). Tissue recovery of essential oils used as feed additive in piglet feeding and impact on nutrient digestibility. In *Proc. Soc. Nutr. Physiol* (Vol. 15, p. 60).

**Struhsaker, T. T., Cooney, D. O., & Siex, K. S.** (1997). Absorptive capacity of charcoals eaten by Zanzibar red colobus monkeys: its function and its ecological and demographic consequences. *International Journal of Primatology*, 18: 61-72.

**Sweeney, T., & O'Doherty, J. V.** (2016). Marine macroalgal extracts to maintain gut homeostasis in the weaning piglet. *Domestic animal endocrinology*, 56, S84-S89.

**Tohamy, M. M., Shaaban, H. A., Ali, M. A., & Hasanain, A. M.** (2019). Effect of Spirulina platensis as Nutrition Source on the Chemical, Rheological and Sensory Properties of Spreadable Processed Cheese. *Journal of Biological Sciences*, 19: 84-91.

**Windisch, W. M., Schedle, K., Plitzner, C., & Kroismayr, A.** (2007). Use of phytogenic products as feed additives for swine and poultry. *J. Anim. Sci.*, 2007, 1-29.

**Zapryanova-Boeva, I. A.** (2011). Effect of plant extracts on the productivity traits in pigs. *Thesis*, 133p. (bg).