



ISOLATION AND CHARACTERISATION OF SHIGA TOXINS PRODUCING *ESCHERICHIA COLI* IN DAIRY COWS IN THE GOVERNORATE OF BLIDA (ALGERIA)

D. BAAZIZE-AMMI¹, S. KECHIH-BOUNAR², A. S. DECHICHA¹,
S. KEBBAL¹, I. GHARBI¹, N. HEZIL¹, Y. CHEBLOUNE³ & D. GUETARNI⁴

¹Institute of Veterinary Sciences, Blida University 1, Algeria; ²Regional Veterinary Laboratory of Draa Ben Khedda, Wilaya of Tizi Ouzou, Algeria; ³Pathogenesis and Lentivirus Vaccination Laboratory, PAVAL Lab., University Grenoble Alps, Grenoble Cedex 9, France; ⁴Biology Department, Faculty of Nature and Life Sciences, Blida University 1, Algeria.

Summary

Baazize-Ammi, D., S. Kechih-Bounar, A. S. Dechicha, S. Kebbal, I. Gharbi, N. Hezil, Y. Chebloune & D. Guetarni, 2024. Isolation and characterisation of Shiga toxins producing *Escherichia coli* in dairy cows in the governorate of Blida (Algeria). *Bulg. J. Vet. Med.*, **27**, No 2, 196–205.

The Shiga toxin producing *Escherichia coli* (STEC) are considered to be one of the most important groups of emerging public health pathogens with cattle being the main reservoir. The objective of this study was to isolate and characterise *Escherichia coli* Shiga toxins in dairy cattle farms. A total of 252 faeces samples were collected from healthy cows belonging to 37 farms. PCR screening of samples for the common sequences of *stx1/stx2* genes and *stx1* and *stx2* genes resulted in a STEC faecal excretion prevalence of 59.5% at the farm level and 26.6% at the individual level. Among positive animals, 85.1% carried STEC with a single *stx1* gene and 14.9% with the *stx1* and *stx2* genes. Immunomagnetic separation was performed on 40 PCR-positive samples (10/10 positive for the *stx1* and *stx2* genes and 30/57 positive only for *stx1*). Biochemical identification revealed the presence of 66 *E. coli* strains (27.5%). The search for virulence genes on these strains by PCR showed that only twenty-two (33.33%) were STEC. The presence of the *stx1*, *stx2*, *ehx* and *eae* genes was characterised in 30.3%, 4.54%, 13.63% and 1.51% of the strains, respectively, indicating that the virulotype with *stx* alone was dominant. Serological identification showed the absence of O157 sero-groups and the presence of O1(2), O2, O18(2), O128 sero-groups. The susceptibility testing of STEC showed 68.18% resistance to chloramphenicol, 63.64% to neomycin, 59.1% to ampicillin, 22.73% to trimethoprim + sulfamethoxazole and 9.1% to amoxicillin + clavulanic acid and nalidixic acid. Four strains showed multi-resistance. Bovine carriage of STEC constitutes a public health risk by contamination of milk and meat. To protect human health, it is necessary to limit the bovine STEC shedding.

Key words: Algeria, cattle, faeces, genes, Shiga toxin producing *Escherichia coli*, virulence

INTRODUCTION

Shiga-toxin-producing *Escherichia coli* (STEC) have been implicated in several worldwide epidemics and sporadic cases resulting from the consumption of contaminated animal foodstuff (Werber *et al.*, 2007). However, other modes of transmission of these pathogens to humans are increasingly reported (Blackburn *et al.*, 2004; Rangel *et al.*, 2005). Non-O157 STEC infections are more common than O157:H7 serotype infections and account for 80% of gastroenteritis as indicated by the national surveillance database in Germany (Werber *et al.*, 2008). The pathogenicity of STEC is attributed to the *stx1* and *stx2* genes which encode Shiga toxins 1 and 2, the *eae* gene coding for the virulence factor intimin and the *ehx* gene encoding the enterohaemolysin A. The combination of these genes is known to increase the risk and severity of infections (Boerlin *et al.*, 1999). STECs have been isolated from a large number of animal species. However, in cattle their presence and excretion are mostly asymptomatic.

Although data on the presence of STEC in Africa are very limited (Effler *et al.*, 2001; Raji *et al.*, 2006), *E. coli* O157 or other STECs have been found in patients and foodstuff from more than twenty countries (Magwira *et al.*, 2005; Al-Gallas *et al.*, 2007; Badri *et al.*, 2009). In Algeria, very few studies have been performed to detect the presence of these bacteria in foodstuffs such as meat (Chahed *et al.*, 2006; Barka *et al.*, 2014). It is therefore essential to identify bovine carriers of these pathogens because they are considered as the natural reservoir.

In this work, the prevalence of Shiga toxin-producing *Escherichia coli* (STEC) in dairy cattle herds from farms located in the governorate of Blida, North Algeria was studied. To this end, evaluation of asymptomatic STEC carriers was deter-

mined by PCR on faecal samples. Positive samples from the first screen were further characterised in terms of STEC strains and virulotypes. The sensitivity to antibiotics of these strains was also analysed.

MATERIALS AND METHODS

Samples collection

The present study involved 252 faecal samples collected from the rectum of healthy adult dairy cows belonging to 37 farms in the governorate of Blida, Algeria. Samples were packaged in sterile 100 mL jars and transported under cold conditions to the laboratory of the Institute of Veterinary Sciences (Blida University 1) for processing.

*Screening for the *stx1* and *stx2* genes by PCR on feces*

The adopted approach was based on the detection of common sequence of *stx1/stx2* genes as a screening test, subsequently followed by seeking the specific sequences of *stx1* and *stx2* genes on positive samples. Sample processing, DNA extraction as well as PCR detection of the common sequence and the *stx1* and *stx2* genes were performed according to the protocol described by Baazize-Ammi *et al.* (2015). The technical conditions of DNA extraction and PCR analysis were first performed at the PAVAL Laboratory, Grenoble, and then transferred to the Viral Respiratory Unit of the Pasteur Institute, Algeria.

Immunomagnetic separation and biochemical identification of strains

From the PCR positive specimens, 40 samples (10 *stx1* and *stx2* positive samples and 30 samples positive only for *stx1*) were selected for isolation and cha-

racterisation of *E. coli* O157. This was done by immunomagnetic separation. Briefly, 25 g of faeces were added to 225 mL of Trypticase-soy broth, homogenised in a Stomacher® 400 Lab Blender (Seward) and incubated at 37 °C for 6 hours for enrichment. The samples thus treated were used for immunomagnetic separation using Dynabeads anti *E. coli* O157 (Invitrogen Dynal, Norway) according to the manufacturer's instructions. The immuno-concentrates obtained were spread on sorbitol MacConkey agar plates and incubated at 37 °C for 24 hours. From the culture plates six characteristic sorbitol-positive colonies and six negative sorbitol colonies were selected and purified for biochemical identification by API 20^E strips (BioMérieux, France).

Screening for *stx1* and *stx2* genes on *E. coli* strain

Using these *E. coli* strains, virulence factors (*stx1*, *stx2*, *eae* and *ehx*) were sought by PCR. DNA extraction was performed with InstaGene™ Matrix reagent (BIO-RAD) according to manufacturer's instructions. Then, GoTaqFlexi DNA Polymerase (Promega, USA) and oligonucleotide primers (Eurogentec, France) were used (Table 1). The PCR mix was prepared as described by Baazize-Ammi *et al.* (2015). The amplification was car-

ried out in a Mastercycler gradient thermocycler (Eppendorf, Germany) according to the following programme: initial denaturation at 94 °C for 3 min, 35 cycles (each comprising a denaturation at 94 °C for 90 s, hybridisation at 59 °C for 90 s and elongation at 72 °C for 120 s), followed by a final elongation at 72 °C for 5 min (Baazize-Ammi *et al.*, 2015).

Serological identification of STEC strains

Serological identification was performed on *E. coli* Shiga toxin strains using the *E. coli* O157 Latex kit (Diagnostic Reagents Oxoid) and Seroscreen DR0300M dry spot kit (Oxoid), according to manufacturer's instructions. The other serogroups were demonstrated using monovalent and polyvalent sera (Eurobio, Sifin) anti somatic O antigens by performing a slide agglutination.

Study of STEC strains sensitivity

For the study of the susceptibility of the STEC strains, a range of eleven antibiotics packaged in the form of discs with a diameter of 9 mm (Oxoid) with the charges recommended by the Algerian Network of the Surveillance of the Resistance of Bacteria to Antibiotics (2011) was used. The antibiogram was performed using the Mueller Hinton agar diffusion method.

Table 1. Nucleotide sequences of primers used

Virulence factors	Primer code	Sequence (5'-3')	bp	References
<i>stx1</i>	B54	AGAGCGATGTTACGGTTTG	388	China <i>et al.</i> (1996), Carneiro <i>et al.</i> (2006)
	B55	TTGCCCCCAGAGTGGATG		
<i>stx2</i>	B56	TGGGTTTTTCTTCGGTATC	807	
	B57	GACATTCTGGACTCTCTT		
<i>eae</i>	B52	AGGCTTCGTCACAGTTG	570	Fremaux <i>et al.</i> (2006)
	B53	CCATCGTCACCAGAGGA		
<i>ehx</i>	hly AF	GCATCATCAAGCGTACGTTCC	534	
	hly AR	AATGAGCCAAGCTGGTTAAGCT		

RESULTS

The detection of the common sequence of *stx1/stx2* genes, based on a PCR screening method, revealed that 67 cows were positive, giving an overall positive rate of 26.6% in individuals and a rate of 59.5% in cattle farms. The separate detection of either *stx1* or *stx2* genes among the positive samples demonstrated that 57 cows carried STECs with the *stx1* gene alone (85.1%) corresponding to 81.8% at the breeding level and 10 cows carrying STECs having simultaneously the *stx1* and *stx2* genes – 14.9% corresponding to 18.2% at the farm level (Table 2).

Biochemical characterisation on API 20^E strips of colonies from Sorbitol Mac-Conkey agar plates allowed identification of 66 *E. coli* strains. The PCR analysis of identified strains for the virulence factors *stx1*, *stx2*, *eae* and *ehx* revealed that only 22 strains (33.33%) were STEC positive because of carrying *stx1* and/or *stx2* genes (Fig. 1). Interpretation of the combined results of virulence factors yielded 5 virulotypes, with a predominance of type 1 (*stx1*⁺ *stx2*⁻ *eae*⁻ *ehx*⁻), followed by type 2 (*stx1*⁺ *stx2*⁻ *eae*⁻ *ehx*⁺) (Table 3).

The serological identification allowed the characterisation of six strains belonging to serogroups O1(2), O2, O18(2),

Table 2. Distribution of *stx1*, *stx1* and *stx2*, and *stx2* genes amongst individual animals and herds

			Common sequence	<i>stx1</i> gene	<i>stx1/stx2</i> genes	<i>stx2</i> gene
Herds	Total (n=37)	(n) (%)	22 59.5			
	Positive (n=22)	(n) (%)		18 81.8	4 18.2	0 0
Animals	Total (n=252)	(n) (%)	67 26.6			
	Positive (n=67)	(n) (%)		57 85.1	10 14.9	0 0




Fig. 1. Electrophoretic analysis of representative PCR products. PCR products (15 µL) from each amplified sample, together with that of the positive control (EDL933), were loaded in wells of 1.5% agarose gel in parallel with a 1000 bp molecular weight DNA marker (MT) and separated by electrophoresis. **A.** Electrophoretic profiles showing the 388 bp product for *stx1* gene (lanes 28-51: *stx1* positive samples; **B.** the 807 bp product for *stx2* (lanes 36 and 49: *stx2* positive samples).

O128 as well as 16 other nonagglutinable isolates (Table 3).

The results of the antibiotic susceptibility study of the 22 STEC strains indicated resistance in 68.18% of strains to chloramphenicol, in 63.64% to neomycin, in 59.1% to ampicillin, in 22.73% to trimethoprim with sulfamethoxazole, and in 9.1% amoxicillin+clavulanic acid and nalidixic acid. Furthermore, multi-resistance against two and more antibiotics at a time, was detected for four strains (18.18%) (Table 4).

DISCUSSION

The PCR search for the common sequence of *stx1/stx2* genes in 252 faecal samples revealed an individual prevalence rate of 26.6%. From these, 85.1% carried only the *stx1* gene while 14.9% had both *stx1* and *stx2* genes, for an overall rate of 59.5% for herds. The presence of STEC in cattle has been reported in many countries, the data are numerous and vary according to individual studies. The numerous observed differences derive from the choice of farms, breed of animals, sampling frequency and methods, and types

Table 3. Combined results for serogroups and virulence factors in 22 STEC strains

Serogroups	Virulence factors (by PCR)				Virulotype
	<i>stx1</i>	<i>stx2</i>	<i>eae</i>	<i>ehx</i>	
N.A.	+	—	—	—	1
N.A.	+	—	—	+	2
N.A.	+	—	—	—	1
N.A.	+	—	—	+	2
O:1	+	—	—	+	2
O:18	+	—	—	—	1
N.A.	+	—	—	+	2
N.A.	+	—	—	+	2
N.A.	+	—	—	—	1
N.A.	+	—	—	—	1
O:1	+	—	—	—	1
N.A.	+	—	—	+	2
O:128	+	—	—	+	2
N.A.	+	—	—	—	1
O:18	+	—	—	—	1
N.A.	+	—	—	—	1
O:2	—	+	—	+	5
N.A.	+	+	—	—	3
N.A.	—	+	+	—	4
N.A.	+	—	—	—	1
N.A.	+	—	—	—	1
N.A.	+	—	—	+	2

N.A.: non agglutinable; Virulotypes: (*stx1*⁺*stx2*[—]*eae*[—]*ehx*[—])=1; (*stx1*⁺*stx2*[—]*eae*[—]*ehx*⁺)=2; (*stx1*⁺*stx2*⁺*eae*[—]*ehx*[—])=3; (*stx1*[—]*stx2*⁺*eae*⁺*ehx*[—])=4 and (*stx1*[—]*stx2*⁺*eae*[—]*ehx*⁺)=5.

Table 4. Strain distribution according to their susceptibility and resistance

Antibiotics	Strain			
	Susceptibility		Resistance	
	n	%	n	%
Ampicillin	9	40.90	13	59.1
Amoxicillin + clavulanic acid	20	90.90	2	9.1
Ceftiofur	22	100	0	0
Neomycin	8	36.36	14	63.64
Gentamicin	22	100	0	0
Trimethoprim + sulfamethoxazole	17	77.27	5	22.73
Nalidixic acid	20	90.90	2	9.1
Enrofloxacin	22	100	0	0
Colistin	22	100	0	0
Chloramphenicol	7	31.82	15	68.18
Nitrofurantoin	22	100	0	0

of analysis (Islam et al., 2008; Rivera et al., 2012).

Isolation and characterisation revealed only 22 STEC strains. For samples tested positive for *stx* genes by PCR and for which STEC could not be isolated, different hypotheses can be proposed: the *stx* element detected by PCR could come from free DNA resulting from bacterial lysis, bacteria found in a viable non-cultivable form (Colwell et al., 1985), or bacteriophages carrying the *stx* gene (Schmidt et al., 2001). This low STEC level could be also explained by the probable loss of *stx* genes occurring during strain isolation. This phenomenon has been reported by Karch et al. (1992) for non-O157 EHEC strains and by Feng et al. (2001) for EHEC O157 strains.

In this study, predominance of the *stx1* gene, followed by the *ehx* gene was observed, while the *stx2* and *eae* genes occurred at lower levels. These results are in line with the work of Scotland et al. (1990) and Willshaw et al. (1992) who reported that production of *stx1* toxin alone was much more common in non-

O157:H7 *E. coli* strains. Similar findings were observed by Tayzar et al. (2013), while higher rates of *stx2* gene were reported by Fernandez et al. (2010), Bosilevac & Koohmaraie (2011) and Ballem et al. (2020). It is important to recall that the *stx2* toxin is more cytotoxic than *stx1* and that *stx2* is associated with high virulence in humans (Rasooly & Do, 2010).

The *eae* gene allowing the attachment and erasure of intestinal cells, fundamental prerequisite for the expression of full virulence of EHEC strains, was carried by one strain only (1.51%). This low rate observed does not necessarily indicate that these strains are not pathogenic for humans. Indeed, atypical strains (absence of the *eae* gene) have already been implicated in haemolytic-uremic syndrome cases in many countries (Werber et al., 2007).

Concerning the *ehx* gene, our results revealed a rate of only 13.63%. According to Sandhu et al. (1996), enterohaemolysin is produced by 98% of non-O157 *eae* positive *E. coli* and by only 36% of *eae* negative strains isolated from cattle, sug-

gesting a high correlation between attachment factor (intimin) and the production of enterohaemolysin (Ehx). Altogether, our data suggest that all isolated STEC strains carry only two genes at most, with a predominance of the *stx* virulotype alone. Indeed, according to Döpfer et al. (2012), this virulotype is found at a higher rate than other virulotypes containing another factor (*eae* and *ehx*) in combination with the *stx* gene. According to Thierry et al. (2020), a total of 10 different virulence patterns were recovered amongst animal species.

Serotyping of the strains showed the absence of the serogroup O157, which could be explained by the intermittent or punctual nature of the excretion and by a seasonal effect. It should be noted that a high prevalence has been reported during summer (Garber et al., 1999). As far as this study was concerned, almost all samples were taken during autumn, winter and spring, from October until April. In dairy farming, the proportion of animals excreting STEC O157 could be estimated at 0.2 to 48.8% of animals (Hussein & Sakuma, 2005). For the other identified serogroups (O1, O2, O18 and O128), they reflect the wide variety of serotypes characterised in cattle, such as 77 different serotypes in dairy cattle faeces and 250 in beef cattle faeces (Hussein & Sakuma, 2005; Hussein & Bollinger, 2005).

The results of the sensitivity study of STEC strains to antibiotics showed resistance to chloramphenicol 68.18%, neomycin 63.64%, ampicillin 59.1%, trimethoprim+sulfamethoxazole 22.73%, amoxicillin+clavulanic acid and nalidixic acid 9.1%. This resistance to antibiotics has been reported by many authors for STEC isolated from various sources (Cergole-Novella et al., 2011; Hossain et al., 2011; Rehman et al., 2013).

CONCLUSION

Based on the results obtained, it may be affirmed that dairy cattle in the governorate of Blida are a reservoir of STEC. It is probable that certain strains isolated during this study are potentially pathogenic for humans or may become so. These strains can be transmitted to humans through direct contact with carrier animals and through raw milk. Waste from these animals can also contaminate the environment and agricultural production through the use of manure.

REFERENCES

- Al-Gallas, N., O. Bahri, A. Bouratbeen, A.B. Haasen & B.A. Ridha, 2007. Etiology of acute diarrhea in children and adults in Tunis, Tunisia, with emphasis on diarrheagenic *Escherichia coli*: Prevalence, phenotyping, and molecular epidemiology. *American Journal of Tropical Medicine and Hygiene*, **77**, 571–582.
- Baazize-Amami, D., O. Gassem, F. Derrar, K. Izri, M. Brahim-Errahmani, J. Gagnon, D. Guetarni, & Y. Chebloune, 2015. Prevalence of asymptomatic carriers of Shiga toxin-producing *Escherichia coli* (STEC) in dairy cattle farms in the governorate of Blida (Algeria). *Bulletin of the Veterinary Institute in Pulawy*, **59**, 23–28.
- Badri, S., A. Fassouane, I. Filliol, M. Hassar & N. Cohen, 2009. Molecular typing of *Escherichia coli* strains isolated from food in Casablanca (Morocco). *Cellular and Molecular Biology*, **55**, 1132–1137.
- Ballem A., S. Gonçalves, I. Garcia-Meniño, S. C. Flament-Simon, J. E. Blanco, C. Fernandes, M. J. Saavedra, C. Pinto, H. Oliveira, J. Blanco, G. Almeida & C. Almeida, 2020. Prevalence and serotypes of Shiga toxin-producing *Escherichia coli* (STEC) in dairy cattle from Northern Portugal. *PLOS One*, 1–12, <https://doi.org/10.1371/journal.pone.0244713>.

- Barka, M. S., N. E. Benyoub, J. E. Henni, B. Boudilmi & M. Kihal, 2014. Genetic characterization of shiga toxin-producing *Escherichia coli* strains isolated from frozen bovine meat in Algeria. *Advances in Environmental Biology*, **8**, 6–13.
- Blackburn, B.G., G. F. Craun, J. S. Yoder, V. Hill, R. L. Calderon, N. Chen, S. H. Lee, D. A. Levy & M. J. Beach, 2004. Surveillance for waterborne-disease outbreaks associated with drinking water – United States, 2001–2002. *Morbidity and Mortality Weekly Report Surveillance Summaries*, **53**, 23–45.
- Boerlin, P., S. A. McEwen, F. Boerlin-Petzold, J. B. Wilson, R. P. Johnson & C. L. Gyles, 1999. Associations between virulence factors of Shiga toxin-producing *Escherichia coli* and disease in humans. *Journal of Clinical Microbiology*, **37**, 497–503.
- Bosilevac, J. M. & M. Koohmaraie, 2011. Prevalence and characterization of non-O157 Shiga toxin-producing *Escherichia coli* isolated from commercial ground beef in the United States. *Applied and Environmental Microbiology*, **77**, 2103–2112.
- Carneiro, L. A., M. C. Lins, F. R. Garcia, A. P. Silva, P. M. Mauller, G. B. Alves, A. C. Rosa, J. R. Andrade, A. C. Freitas-Almeida & M. L. Queiroz, 2006. Phenotypic and genotypic characterisation of *Escherichia coli* strains serogrouped as enteropathogenic *E. coli* (EPEC) isolated from pasteurised milk. *International Journal of Food Microbiology*, **108**, 15–21.
- Cergole-Novella, M. C., A. C. Pignatari, M. Castanheira & B. E. Guth, 2011. Molecular typing of antimicrobial-resistant Shiga-toxin-producing *Escherichia coli* strains (STEC) in Brazil. *Research in Microbiology*, **162**, 117–123.
- Chahed, A., B. China, J. Mainil & G. Daube, 2006. Prevalence of enterohaemorrhagic *Escherichia coli* from serotype O157 and other attaching and effacing *Escherichia coli* on bovine carcasses in Algeria. *Journal of Applied Microbiology*, **101**, 361–368.
- China, B., V. Pirson & J. Mainil, 1996. Typing of bovine attaching and effacing *Escherichia coli* by multiplex *in vitro* amplification of virulence-associated genes. *Applied and Environmental Microbiology*, **62**, 3462–3465.
- Colwell, R. R., P. R. Brayton, D. J. Grimes, D. R. Roszak, S. A. Huq & L. M. Palmer, 1985. Viable, but non-culturable *Vibrio cholerae* and related pathogens in the environment: Implications for release genetically engineered microorganisms. *Bio/Technology*, **3**, 817–820.
- Döpfer, D., L. Geue, S. Schares, B. Mintel, B. Hoffmann & E. A. J. Fischer, 2012. Dynamics of shiga-toxin producing *Escherichia coli* (STEC) and their virulence factors in cattle. *Preventive Veterinary Medicine*, **103**, 22–30.
- Effler, E., M. Isaacson, L. Arntzen, R. Heenan, P. Canter, T. Barrett, L. Lee, C. Mambo, W. Levine, A. Zaidi & P. M. Griffin, 2001. Factors contributing to the emergence of *Escherichia coli* O157 in Africa. *Emerging Infectious Diseases*, **7**, 812–819.
- Feng, P., M. Dey, A. Abe & T. Takeda, 2001. Isogenic strain of *Escherichia coli* O157:H7 that has lost both Shiga toxin 1 and 2 genes. *Clinical and Diagnostic Laboratory Immunology*, **8**, 711–717.
- Fernandez, D., K. Irino, M. E. Sanz, N. L. Padola & A. E. Parma, 2010. Characterization of Shiga toxin-producing *Escherichia coli* isolated from dairy cows in Argentina. *Letters in Applied Microbiology*, **51**, 377–382.
- Fremaux, B., S. Raynaud, L. Beutin & C. Vernozy-Rozand, 2006. Dissemination and persistence of Shiga toxin-producing *Escherichia coli* (STEC) strains on French dairy farms. *Veterinary Microbiology*, **117**, 180–191.
- Garber, L., S. Wells, L. Schroeder-Tucker & K. Ferris, 1999. Factors associated with fecal shedding of verotoxin-producing *Escherichia coli* O157 on dairy farms. *Journal of Food Protection*, **62**, 307–312.
- Hossain, Z., P. Sultana, S. Deb & M. M. Ahmed, 2011. Multidrug resistance in large-

- plasmid-associated presumptive enterohaemorrhagic *Escherichia coli* isolated from contaminated lake water. *Bangladesh Journal of Microbiology*, **28**, 33–40.
- Hussein, H. S. & L. M. Bollinger, 2005. Prevalence of Shiga toxin-producing *Escherichia coli* in beef cattle. *Journal of Food Protection*, **68**, 2224–2241.
- Hussein, H. S. & T. Sakuma, 2005. Prevalence of Shiga toxin-producing *Escherichia coli* in dairy cattle and their products. *Journal of Dairy Science*, **88**, 450–465.
- Islam, M. A., A. S. Mondol, E. De Boer, R. R. Beumer, M. H. Zwietering, K. A. Talukder, A. E. Heuvelink, 2008. Prevalence and genetic characterization of Shiga toxin-producing *Escherichia coli* isolates from slaughtered animals in Bangladesh. *Applied and Environmental Microbiology*, **74**, 5414–5421.
- Karch, H., T. Meyer, H. Rüssmann & J. Heesemann, 1992. Frequent loss of Shiga-like toxin genes in clinical isolates of *Escherichia coli* upon subcultivation. *Infection and Immunity*, **60**, 3464–3467.
- Magwira, C. A., B. A. Gashe & E. K. Collison, 2005. Prevalence and antibiotic resistance profiles of *Escherichia coli* O157:H7 in beef products from retail outlets in Gaborone, Botswana. *Journal of Food Protection*, **68**, 403–406.
- Raji, M. A., U. Minga & R. Machangu, 2006. Current epidemiological status of enterohaemorrhagic *Escherichia coli* O157:H7 in Africa. *Chinese Medical Journal*, **119**, 217–222.
- Rangel, J. M., P. H. Sparling, C. Crowe, P. M. Griffin & D. L. Swardlow, 2005. Epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982–2002. *Emerging Infectious Diseases*, **11**, 603–609.
- Rasooly, R. & P. M. Do, 2010. Shiga toxin Stx2 is heat stable and not inactivated by pasteurization. *International Journal of Food Microbiology*, **136**, 290–294.
- Rehman, M. U., M. Rashid, J. Ahmad Sheikh, S. Ahmad Wani & S. Farooq, 2013. Multi-drug resistance among Shiga toxin producing *Escherichia coli* isolated from bovines and their handlers in Jammu region, India. *Veterinary World*, **6**, 655–658.
- Réseau Algérien de la surveillance de la résistance des bactéries aux antibiotiques, 2011. Standardisation de l'antibiogramme à l'échelle nationale (médecine humaine et vétérinaire). 6^{ème} édition, <https://docplayer.fr/27601032-Standardisation-de-l-antibiogramme-a-l-echelle-nationale.html> (08 July 2022, date last accessed).
- Rivera, F. P., E. Sotelo, I. Morales, F. Menacho, A.M. Medina, R. Evaristo, R. Valencia, L. Carbajal, J. Ruiz & T. J. Ochoa, 2012. Detection of Shiga toxin-producing *Escherichia coli* (STEC) in healthy cattle and pigs in Lima, Peru. *Journal of Dairy Science*, **95**, 1166–1169.
- Sandhu, K. S., R. C. Clarke, K. McFadden, A. Brouwer, M. Louie, J. Wilson, H. Lior & C. L. Gyles, 1996. Prevalence of *eaeA* gene in virotoxicogenic *Escherichia coli* strains from dairy cattle in southwest Ontario. *Epidemiology & Infection*, **116**, 1–7.
- Schmidt, H., W. L. Zhang, U. Hemmrich, S. Jelacic, W. Brunder, P. I. Tarr, U. Do-brindt, J. Hacker & H. Karch, 2001. Identification and characterization of a novel genomic island integrated at *selC* in locus of enterocyte effacement-negative, Shiga toxin-producing *Escherichia coli*. *Infection and Immunity*, **69**, 6863–6873.
- Scotland, S. M., G. A. Willshaw, H. R. Smith & B. Rowe, 1990. Properties of strains of *Escherichia coli* O26:H11 in relation to their enteropathogenic or enterohemorrhagic classification. *Journal Infectious Diseases*, **162**, 1069–1074.
- Tayzar, A. C, A. A. Saleha, A. M. Rahim, M. Murugaiyah & A. H. Shah, 2013. Occurrence of non-O157 Shiga toxin-producing *Escherichia coli* in healthy cattle and goats and distribution of virulence genes among isolates. *African Journal of Microbiology Research*, **7**, 1703–1707.
- Thierry, S. I. L., J. E. Gannon, Y. Jaufeerally-Fakim, S. J. Santchurn, 2020. Shiga-toxigenic *Escherichia coli* from animal food sources in Mauritius: Prevalence, serogroup diversity and virulence profiles.

International Journal of Food Microbiology, **324**, 108589.

Journal of Infectious Diseases, **166**, 797–802.

Werber, D., S. C. Behnke, A. Fruth, R. Merle, S. Menzler, S. Glaser, L. Kreienbrock, R. Prager, H. Tschape, P. Roggentin, J. Bockemuhl & A. Ammon, 2007. Shiga toxin-producing *Escherichia coli* infection in Germany: Different risk factors for different age groups. *American Journal of Epidemiology*, **165**, 425–434.

Paper received 17.03.2022; accepted for publication 30.06.2022

Werber, D., L. Beutin, R. Pichner, K. Stark & A. Fruth, 2008. Shiga Toxin-producing *Escherichia coli* serogroups in food and patients, Germany. *Emerging Infectious Diseases*, **14**, 1803–1806.

Correspondence:

Willshaw, G. A., S. M. Scotland, H. R. Smith & B. Rowe, 1992. Properties of vero Cytotoxin-producing *Escherichia coli* of human origin of O serogroups other than O157.

Djamila Baazize-Ammi
Institute of Veterinary Sciences,
Blida University 1, Algeria
email: baazizedjamila@yahoo.fr